

New equation on the low dimensional Calabi–Yau metrics*

Dmitry Egorov

Abstract

In this paper we introduce a new equation on the compact Kähler manifolds. Solution of this equation corresponds to the Calabi–Yau metric. New equation differs from the Monge–Ampère equation considered by Calabi and Yau.

1 Introduction

Let (M, ω) be the compact Kähler n -manifold. Calabi made the following conjecture proven by Yau: If $c_1(M) = 0$, then there exists a Riemannian metric with the holonomy group contained in $SU(n)$ [1, 2]. Explicitly, Calabi conjecture is stated as follows: There exists unique real function φ such that:

$$(\omega + i\partial\bar{\partial}\varphi)^n = e^F \omega^n, \quad \int_M \varphi \omega^n = 0, \quad (1)$$

where F is a smooth real function on M such that $\int_M e^F \omega^n = \int_M \omega^n$. It is assumed that $\omega + i\partial\bar{\partial}\varphi > 0$.

Existence of the solution for (1) implies existence of the Ricci-flat metric on M .

Note that the Monge–Ampère equation describes the deformation of the Kähler form. We study the deformation of the holomorphic volume form on the Kähler manifold M in complex dimensions 2 and 3.

*This work was supported in part by Russian Foundation for Basic Research (grant 09-01-00598-a) and the Council of the Russian Federation Presidential Grants (projects NSh-7256.2010.1 and MK-842.2011.1).

Theorem. *Let M be a compact Kähler n -manifold. Suppose $c_1(M) = 0$. Denote holomorphic volume form by Ω . For $n = 2, 3$ there exists a solution of the following equation:*

$$(\Omega + dd^s\psi) \wedge (\bar{\Omega} + dd^s\bar{\psi}) = e^F \Omega \wedge \bar{\Omega}, \quad (2)$$

where ψ is a complex n -form such that $\tilde{\Omega} = \Omega + dd^s\psi$ is a stable primitive form; F is a smooth real function such that $\int_M e^F \Omega \wedge \bar{\Omega} = \int_M \Omega \wedge \bar{\Omega}$. Here d^s is the symplectic differential operator defined in §2.

Here $\tilde{\Omega}$ is said to be stable if there exists some complex structure J on M such that $\tilde{\Omega}$ is a holomorphic volume form with respect to J .

The main theorem implies the existence of the Calabi–Yau metric on M . Since $\tilde{\Omega} = \Omega + dd^s\psi$ is the holomorphic volume form, $\tilde{\Omega} \wedge \bar{\tilde{\Omega}}$ is proportional to the volume form. Then there exists a real function F such that

$$e^F \Omega \wedge \bar{\Omega} = \omega^n,$$

where F is a smooth real function such that $\int_M e^F \Omega \wedge \bar{\Omega} = \int_M \Omega \wedge \bar{\Omega}$. By the main theorem, there exists $\tilde{\Omega}$ such that

$$\tilde{\Omega} \wedge \bar{\tilde{\Omega}} = \omega^n. \quad (3)$$

Recall that n -form is called primitive if

$$\Omega \wedge \omega = 0. \quad (4)$$

Stable forms $\tilde{\Omega}$ and ω are closed and satisfy (3),(4). Therefore they set up an integrable $SU(n)$ -structure. Hence the holonomy group of M is contained in $SU(n)$.

In §2 we recall necessary facts from the symplectic Hodge theory. In §3 we prove the main theorem. The proof follows from Yau’s theorem. In §4 we discuss possible generalizations of the main theorem.

The author is grateful to Iskander Taimanov for drawing attention to this area of mathematics and constant support and to Andrey Mironov for useful conversations.

2 Symplectic Hodge theory

Let (M, ω) be a symplectic manifold of real dimension $2n$. By $*_s$ denote a symplectic Hodge star. The action of $*_s$ on the k -forms is uniquely determined by the following formula:

$$\alpha \wedge *_s \beta = (\omega^{-1})^k(\alpha, \beta) \frac{\omega^n}{n!}.$$

By definition, put

$$d^s = (-1)^{k+1} *_s d *_s. \quad (5)$$

Note that d^s reduces degree of the form by one and $dd^s = -d^s d$. Kähler manifolds satisfy the following condition (dd^s -lemma):

$$\text{im } d \cap \ker d^s = \ker d \cap \text{im } d^s = \text{im } dd^s.$$

Lemma 1. *Let M be a Kähler n -manifold. Suppose Ω and $\tilde{\Omega}$ are primitive cohomologous n -forms. Then $\tilde{\Omega} = \Omega + dd^s \psi$, where ψ is some n -form.*

Proof. The following formula for primitive n -forms [5] (see also [6]):

$$*_s \eta = (-1)^{\frac{n(n+1)}{2}} \eta \quad (6)$$

implies that d -closed forms are d^s -closed as well. By the dd^s -lemma, $\tilde{\Omega} - \Omega$ takes the form $dd^s \psi$. $\square$

3 Proof of the main theorem

Recall the definition of a stable form by Hitchin [3, 4]. Let V be a real m -dimensional space. A real form $\rho \in \Lambda^p V^*$ is called stable if the orbit of ρ under the natural action of $GL(V)$ is open.

If $p = 2$ and $m = 2n$, then we obtain the non-degeneracy condition of the symplectic form. Recall that any compact Kähler 2-manifold with $c_1(M) = 0$ is hyper-Kähler. Then by dimensional reasons real and imaginary parts of the holomorphic volume form are symplectic and hence are stable.

If $p = 3$ and $m = 6$, then by definition stable forms are real and imaginary parts of the holomorphic volume form.

We need the Moser theorem for oriented manifolds [7].

Theorem. *Let M be an oriented manifold. Suppose there exists a family of cohomologous symplectic forms $\{\omega_t\}, t \in [0, 1]$. Then there exists an orientation preserving diffeomorphism ϕ_t such that*

$$\phi_t^* \omega_t = \omega_0,$$

and ϕ_0 is the identity map.

Proof. By Yau's theorem, there exists $\tilde{\omega} \sim \omega$ compatible with Ω .

In his proof Yau uses the continuity method. Therefore we get a family of cohomologous symplectic forms such that

$$\omega_0 = \omega, \quad \omega_1 = \tilde{\omega}.$$

Denote ϕ_1 by ϕ . We claim that $\phi^* \Omega$ is the required $\tilde{\Omega}$. Indeed, act by ϕ^* on the compatible pair: Ω and $\tilde{\omega} = (\phi^{-1})^* \omega$. Diffeomorphism preserves compatibility conditions (3),(4) and stability. Since ϕ is isotopic to the identity map, $\tilde{\Omega} = \phi^* \Omega \in [\Omega]$.

Stable forms $\tilde{\Omega}$ and ω are closed and compatible in the sense of (3),(4). Hence they define a Riemannian metric with holonomy group contained in $SU(n)$.

Let's prove that $\tilde{\Omega}$ is a solution of (2). Since $\tilde{\Omega}$ is the holomorphic volume form with respect to some complex structure, $\tilde{\Omega} \wedge \bar{\tilde{\Omega}}$ is proportional to the volume form. Therefore there exists a smooth real function F such that

$$e^F \tilde{\Omega} \wedge \bar{\tilde{\Omega}} = \omega^n, \tag{7}$$

$$\int_M e^F \tilde{\Omega} \wedge \bar{\tilde{\Omega}} = \int_M \tilde{\Omega} \wedge \bar{\tilde{\Omega}}.$$

Substituting (7) in (3), we obtain the required equation (2):

$$\tilde{\Omega} \wedge \bar{\tilde{\Omega}} = \omega^n = e^F \Omega \wedge \bar{\Omega}.$$

□

The key moment of the proof is the stability of the holomorphic volume form. Therefore this proof can not be generalized to higher dimensions.

4 Conclusion

Here we discuss possible extensions of the work.

Let (M, ω) be a symplectic non-Kähler manifold of dimension $2n$. Direct generalization of the Monge–Ampère equation is as follows:

$$(\omega + dJd\varphi)^n = e^F \omega^n, \quad (8)$$

where J is an almost complex structure. If J is integrable, then this equation turns into the Monge–Ampère equation.

Equation (8) doesn’t give anything new. It is proved in [8, 9] that if for any compatible RHS there exists a solution, then J is integrable.

It is interesting to find an analogue of this result for new equation. In generalized complex geometry a non-degenerate 2-form is called an almost symplectic structure. Integrability stands for closedness.

Assume that for any compatible RHS there exists a solution of the equation:

$$\tilde{\Omega} \wedge \bar{\tilde{\Omega}} = e^F \Omega \wedge \bar{\Omega}, \quad \tilde{\Omega} = \Omega + dd^s \psi.$$

Does it imply integrability, i.e., closedness, of the almost symplectic structure?

Let’s introduce the second possible extension of the current work. The Monge–Ampère equation is generalized to non-Kähler symplectic 4-manifolds by Donaldson–Yau–Weinkove–Tosatti. It is suggested in [10] to consider the following equation:

$$\tilde{\omega}^n = e^F \omega^n, \quad (9)$$

where $\tilde{\omega} \sim \omega$ are symplectic forms. This equation is called the Calabi–Yau equation.

In [11, 12, 13] authors assume that the curvature of the Chern connection satisfies some restrictions. Then they prove the existence of the unique solution for (9).

Consider a generalization of the new equation to non-Kähler manifolds. In our opinion the non-Kähler Hermitian manifolds with trivial canonical bundle (see for example [14]) are natural candidates for generalization of the existence theorem.

Finally, in the Kähler case it is unclear: Does there exist generalization of the existence theorem for $n > 3$?

References

- [1] E. Calabi, On Kähler manifolds with vanishing canonical class, Algebraic geometry and topology. A symposium in honor of S. Lefschetz, pp. 78–89. Princeton University Press, Princeton, N.J., 1957
- [2] S.-T. Yau, On the Ricci curvature of compact Kähler manifold and the complex Monge–Ampère equation I, Comm. on pure and appl. math. V. 31, 339–411, 1978
- [3] N.J.Hitchin, The geometry of three-forms in six dimensions, J. Differential Geometry 55 (2000), 547–576.
- [4] N.J.Hitchin, Stable forms and special metrics, in "Global Differential Geometry: The Mathematical Legacy of Alfred Gray", M. Fernández and J. A. Wolf (eds.), Contemporary Mathematics 288, American Mathematical Society, Providence (2001).
- [5] A. Weil , Introduction à l'Étude des variété Kahlériennes, Publications de l'Instiut de Mathématique de l'Université de Nancago VI, Hermann, Paris, 1958
- [6] Li-Sheng Tseng and S.-T. Yau , Cohomology and Hodge theory on symplectic manifolds: I, arXiv:0909.5418
- [7] J. Moser, On the volume elements of a manifold, Trans. AMS 120 (1965), 286–294.
- [8] P. Delanoë, Sur l'analogie presque-complexe de l'equation de Calabi–Yau, Osaka J. Math. 33 (1996), 829–846.
- [9] Wang Hongyu, Zhu Peng. On a generalized Calabi–Yau equation, arXiv:0911.0784
- [10] S.K. Donaldson, Two-forms on four-manifolds and elliptic equations, Inspired by S.S.Chern, World Scientific, 2006
- [11] B. Weinkove, The Calabi–Yau equation on almost Kähler four-manifolds, J.Diff.Geometry, V. 76, N. 2, 317–349, 2007

- [12] V. Tosatti, B. Weinkove, S.-T. Yau, Taming symplectic forms and the Calabi–Yau equation, Proc. London Math. Soc. V. 97, N. 2, 401–424, 2008
- [13] V. Tosatti and B. Weinkove, The Calabi–Yau equation, symplectic forms and almost complex structures, arXiv:0901.1501
- [14] J. Fine, D. Panov, Hyperbolic geometry and non-Kähler manifolds with trivial canonical bundle, arXiv:0905.3237

AMMOSOV NORTHEASTERN FEDERAL UNIVERSITY, YAKUTSK, 677000,
RUSSIA

*e-mail:*egorov.dima@gmail.com